

## New Hampshire Medicaid Fee-for-Service Program

### Atopic Dermatitis Criteria

Approval Date: July 12, 2022

### Medications

Brand Names	Generic Names	Dosage
Cibinqo™	abrocitinib	50 mg, 100 mg, 200 mg tablets
Eucrisa®	crisaborole	2% ointment
Elidel®	pimecrolimus	1% cream
Opzelura™	ruxolitinib	1.5% cream
Protopic®	tacrolimus	0.03%, 0.1% ointment
Adbry™	tralokinumab-ldrm	150 mg/mL prefilled syringe
Rinvoq®	upadacitinib	15 mg, 30 mg tablets

For requests for dupilumab (Dupixent®) use the Dupixent® criteria

### Criteria for Approval

#### Topical Therapy

1. FDA-approved indication and age:
  - a. **Eucrisa® (crisaborole)**: Topical treatment of mild to moderate atopic dermatitis in patients  $\geq$  3 months of age; **OR**
  - b. **Elidel® (pimecrolimus)**: Second-line therapy for the short-term and non-continuous chronic treatment of mild to moderate atopic dermatitis in non-immunocompromised adults and children  $\geq$  2 years old who have failed to respond adequately to other topical prescription treatments or when those treatments are not advisable; **OR**
  - c. **Opzelura™ (ruxolitinib)**: Second-line therapy for the short-term and non-continuous chronic treatment of mild to moderate atopic dermatitis in non-immunocompromised patients  $\geq$  12 years of age who have failed to respond adequately to other topical prescription treatments or when those treatments are not advisable; **OR**
  - d. **Protopic® (tacrolimus)**: Second-line therapy for the short-term and non-continuous chronic treatment of moderate to severe atopic dermatitis in non-immunocompromised adults and

children who have failed to respond adequately to other topical prescription treatments for atopic dermatitis or when those treatments are not advisable

- i. 0.03% ointment approved for patients  $\geq$  2 years old; **OR**
  - ii. 0.1% ointment approved for patients  $\geq$  16 years old; **AND**
2. Patient has a defined failure or contraindication or intolerance to a trial of topical corticosteroids. In general, a trial constitutes two weeks for high-potency topical corticosteroids (e.g., diflorasone diacetate) and four weeks for low-potency topical corticosteroids (e.g., hydrocortisone acetate); **AND**
  3. Opzelura™ only: Patient has a defined failure or contraindication or intolerance to a trial of topical calcineurin inhibitors (e.g., pimecrolimus or tacrolimus) or topical phosphodiesterase-4 inhibitor (e.g., crisaborole); **AND**
  4. Prescribed utilization is for short-term (up to six consecutive weeks at a time) therapy or for non-continuous intermittent therapy (up to one year in duration).

**Length of Approval:** Two months

**Renewal:** Six months

### **Systemic Therapy**

1. FDA-approved indication and age:
  - a. **Cibinqo™ (abrocitinib):** indicated for the treatment of adults with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is inadvisable; **OR**
  - b. **Adbry™ (tralokinumab-ldrm):** indicated for the treatment of moderate-to-severe atopic dermatitis in adult patients whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable; **OR**
  - c. **Rinvoq® (upadacitinib):** indicated for the treatment of adults and pediatric patients  $\geq$  12 years of age with refractory, moderate to severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies is not advisable; **AND**
2. Prescriber is a dermatologist, immunologist, or allergist (or one has been consulted); **AND**
3. Patient is  $\geq$  18 years of age or  $\geq$  12 years of age for Rinvoq®; **AND**
4. Patient has a defined failure, contraindication, or intolerance to a trial of topical corticosteroids. In general, a trial constitutes two weeks for high-potency topical corticosteroids (e.g., diflorasone diacetate) and four weeks for low-potency topical corticosteroids (e.g., hydrocortisone acetate); **AND**

5. Patient has a defined failure, contraindication, or intolerance to a trial of pimecrolimus **OR** a trial of tacrolimus. A trial constitutes at least one month of therapy; **AND**
6. Patient has a defined failure, contraindication, or intolerance to a trial of Eucrisa® (crisaborole). A trial constitutes at least one month of therapy; **AND**
7. Prescribed utilization is for short-term (up to six consecutive weeks at a time) therapy or for non-continuous intermittent therapy (up to one year in duration); **AND**
8. Patient will not receive concurrent therapy with any other monoclonal antibody biologics (e.g., Tezepelumab, omalizumab, mepolizumab, reslizumab, dupilumab).

## Criteria for Denial

1. Failure to meet criteria for approval; **OR**
2. Treatment of psoriasis; **OR**
3. Treatment of infected atopic dermatitis; **OR**
4. Treatment of Netherton’s syndrome.

## References

Available upon request

## Revision History

Reviewed by	Reason for Review	Date Approved
Pharmacy and Therapeutic Committee	New	11/02/2006
Commissioner	New	11/16/2006
DUR Board	Review	10/25/2010
Commissioner	Approval	02/10/2011
DUR Board	Revision	10/24/2017
Commissioner	Approval	12/05/2017
DUR Board	Revision	03/12/2019
Commissioner Designee	Approval	04/05/2019
DUR Board	Revision	10/28/2019
Commissioner Designee	Approval	12/03/2019
DUR Board	Revision	06/30/2020
Commissioner Designee	Approval	08/07/2020
DUR Board	Revision	12/15/2020

Reviewed by	Reason for Review	Date Approved
Commissioner Designee	Approval	02/24/2021
DUR Board	Revision	06/02/2022
Commissioner Designee	Approval	07/12/2022